

Need ideas for your STD Prevention underwear design? Want to learn more about protecting yourself and your partners?

Did you know...

It's normal to be afraid of telling your partners about an STD exposure.
Learn more about [STDs and communication](#)

HPV can cause cervical cancer in women yet infection often has no symptoms.
Learn more about [HPV](#)

The highest rates of gonorrhea infection are among sexually active teens.
Learn more about [gonorrhea](#)

Chlamydia can be transmitted to the throat during oral sex.
Learn more about [chlamydia](#)

Many people infected with syphilis do not have any symptoms for years, yet remain at risk for late complications if they are not treated.
Learn more about [syphilis](#)

Chlamydia, gonorrhea and syphilis can be successfully treated with antibiotics.
Learn more about [STD treatment](#)

Approximately 40,000 people become infected with HIV each year in the U.S.
Learn more about [HIV](#)

One out of five adolescents and adults has genital herpes infection.
Learn more about [genital herpes](#)

Viral hepatitis can be transmitted sexually and is the leading cause of liver transplants.
Learn more about [hepatitis](#)

Having an STD can put you at higher risk for contracting HIV.
Learn more about [HIV](#)

Genital warts, caused by the human papillomavirus (HPV), are spread by skin-to-skin contact from one person to another during sex.
Learn more about [genital warts](#)